arXiv:1212.0435v1 [cond-mat.stat-mech] 3 Dec 2012

Network Growth with Arbitrary Initial Conditions: Analytical Results for Uniform and Preferential Attachment

Babak Fotouhi and Michael G. Rabbat
Department of Electrical and Computer Engineering
McGill University, Montréal, Québec, Canada
Email: `babak.fotouhi@mail.mcgill.ca`, `michael.rabbat@mcgill.ca`

This paper provides exact expressions for the expected degree distribution of a given network that is subject to growth, as a function of time. We consider both uniform attachment, where incoming nodes form links to existing nodes selected uniformly at random, and preferential attachment, when probabilities are assigned proportional to the degrees of the existing nodes. We consider the cases of single and multiple links being formed by each newly-introduced node. The initial conditions are arbitrary, that is, the solution depends on the degree distribution of the initial graph which is the substrate of the growth. Previous work in the literature focuses on the asymptotic state, that is, when the number of nodes added to the initial graph tends to infinity, rendering the effect of the initial graph negligible. Our contribution provides a solution for the expected degree distribution as a function of time, for arbitrary initial condition. Previous results match our results in the asymptotic limit.

I. INTRODUCTION

The complex network literature spans various strands of research such as sociology [1–3], economics [4, 5], computer science [6–8], marketing [9–12], epidemiology [13, 14], genetics [15], and bibliometrics [16]. These domains aim to extract macro-scale behavior from given micro-scale interactions..

The structure of the underlying graph, which connects the agents and consequently regulates their interactions, is necessary for studying the dynamism of various phenomena, such as flow of information (news, rumors, trends, etc.) in the society [17, 18], resilience against node or link failures (for the internet, it means survival of the system if certain nodes are shut down) [19, 20], pace of diffusion of a contagious disease throughout a population and also optimal immunization strategies [21–23], the effect of the network structure among actors on their chance of winning awards [24], to name a few. Models have been proposed to emulate different structural properties observed in real life graphs [25–34].

In many applications, such as the world wide web [35, 36] and scientific collaborations [37], networks are dynamic, that is, subject to growth. This provides motivation to view the problem of network formation dynamically. In this formulation, nodes are introduced successively, and they select from existing nodes whom to attach to. It mimics, for example, the mechanism by which new papers cite existing ones. [34] takes this approach. Also, in [38–41] the problem is tackled by the conventional techniques of polymer physics. Both of these approaches employ approximations to solve the problem. In what follows, we go over these approximations and the corresponding results.

A. Previous Work: Network Growth

In [34], the growth process starts from a small number of m_0 nodes. Then, nodes are introduced one per unit time. Each node picks $m < m_0$ existing nodes to link to, with probabilities assigned to them proportional to their degrees. This means that an existing node with a higher degree will be more likely to attract the newly-introduced node. Denote the degree distribution of the graph when the total number of nodes is N by $p_k(N)$. Note that t and N are interchangeable in this scheme, since nodes are introduced one at a time. Their result can be expresses as follows:

$$\lim_{N\to\infty} p_k(N) \sim k^{-3}. \tag{1}$$

The analysis is done within the mean-field simplification and the solution is valid in the asymptotic case of $t \to \infty$. In [30], this result is ameliorated by reformulating the problem more rigorously, perhaps more congenial to the fastidious reader. Denote by β the number of links that each newly-born node emanates to the existing nodes. The process starts from a $4-$cycle. Let $p_k(N)$ be defined as above. Also, define:

$$\alpha_{\beta,k} \overset{\text{def}}{=} \frac{2(\beta)(\beta+1)}{k(k+1)(k+2)}. \tag{2}$$

The authors show that(Theorem 1 in [30]) in the limit $t \to \infty$, for any positive ϵ and for $0 \leq k \leq N^{1/5}$, the following holds:

$$(1-\epsilon)\alpha_{\beta,k} \leq p_k(N) \leq (1+\epsilon)\alpha_{\beta,k}. \tag{3}$$

Note that the expression in (2) agrees with (1) for large values of k.

The problem is also closely related to the so-called

Polya's urn problem [42] in combinatorics. Given a finite number of bins, additional balls arrive one at a time. With a given probability, a new bin is created for the new ball. The ball otherwise joins an existing bin. It picks the destination bin with probabilities dependent on the existing number of balls within the bins. In [42], the case where probabilities are proportional to m^γ is solved. $m = \gamma$ is akin to the linear preferential attachment scheme mentioned above.

A novel way to tackle the problem was presented in [38–41, 43] by employing the master equation approach which authors borrow from polymer physics. The authors obtain (2) for the case of $\beta = 1$ (equation (2) in[38], (5) in [43], (2) in [39] and (2) in [41]). For a treatment of finite size effects (when N is not infinitely large) with primary focus on nodes with degree $k_{\max} \approx \sqrt{N}$, see [41]. Also, in [38], the uniform attachment scheme is also examined. This means that, new nodes attach to existing nods with equal probabilities, regardless of their degrees. If we start from a single node at the outset, the resulting graph is called a *Random Recursive Tree (RRT).* The result presented in [38] for the asymptotic degree distribution of RRTs is as follows:

$$\lim_{t\to\infty} p_k(t) = \frac{1}{2^k}. \tag{4}$$

The same result is also presented in Theorem 1 in[44] and equation (49) in [45] following a combinatorial approach.

B. Exact Solution, Motivation

Previous work has been primarily revolved around the asymptotic degree distribution, that is, when the number of nodes tends to infinity. Also, in some case, further simplification is acquired by limiting the range of degrees. For long times, the effect of the initial graph is neglected. In this contribution, we start from an initial graph with known degree distribution n_k. We solve for the expected degree distribution at time t. We consider both uniform and linear preferential attachment (eventuating in a scale-free graph in the long run). The exact solution, first develops intuition about the growth process, and the path that the system undergoes until it reaches the steady state. More importantly, the effect of the initial conditions is taken into account. Different substrates reach the equilibrium approximation of the degree distribution which is at hand, with different paces. The exact solution illuminates the effect of the initial condition on the accuracy of the abovementioned approximations.

Equipped with the exact solution, one can also examine the short-time behavior, in marked contrast with the convention, which limits the solution to the long-time behavior. As an example of how the need for extracting the short-time growth of an existing graph is elicited in realistic applications, consider the network of supporters in a political campaign. Nodes are fanatics who absorb new people into the campaign, causing the network to expand throughout the potential electorate. The change in the network of followers in one day is not substantial compared to the existing size of the network. As another example, consider the social network within a country, with a small number of immigrants joining and enlarging the network. The number of immigrants typically constitutes a small fraction of the population of the host country (with possible exceptions of wars or other abrupt phase transitions, to minor degrees). Then, if one wants to study the social network of the host country, the conventional models cease to perform, because the fraction of new nodes to existing nodes does not tend to infinity, but is small. The same is true for any slowly-growing realistic network where the extrapolation of the near future provided information on the current state is called for.

C. Organization of the Paper

First in subsection II A we consider the uniform attachment scheme, with each newly introduced node linking to one existing node picked uniformly at random. We compare our results with the ones present in the literature. Then in subsection II B we consider the uniform attachment for multiple linking, where each new node connects to β existing nodes drawn uniformly at random. In III we examine the preferential attachment scheme. First in III A we consider each new node linking to only one existing node with probabilities assigned to existing nodes proportional to their degrees. Then in III B we assume each new node attaches to β existing nodes. So each new node has degree β upon birth. We solve for the expected degree distribution in all cases. Throughout the paper, we compare our theoretical findings with simulations.

II. UNIFORM ATTACHMENT

A. Single Connection

We start from an initial connected graph at time $t = 0$ with $N(0)$ nodes. We denote the degree distribution at the outset by n_k. At each timestep, a new node is introduced. It picks one of the existing nodes uniformly at random and connects to it. Let α represent the rate at which new nodes are introduced, that is, $\alpha\Delta t$ nodes are added in a time interval of duration Δt. Nodes are added one by one (if the initial condition is a single node, the resulting graph will be the conventional Random Recursive Tree [38, 44, 45]). At time t there are $N(t) = N(0) + \alpha t$ nodes. Let $N_k(t)$ denote the expected number of nodes whose degree is k at time t. Let us focus on the expected variation in $N_k(t)$ in an infinitesimal time increment Δt within which $\alpha \Delta t$ new nodes are added.

With probability $\frac{\alpha N_k(t)}{N(t)}$, a node with degree k receives a link, and its degree increments. Consequently, N_k decrements and N_{k+1} increments, both by one. Similarly, with probability $\frac{\alpha N_{k-1}(t)}{N(t)}$, a node with degree $k-1$ receives a link, hence N_k decrements and N_{k+1} increments, both by one. So we have:

$$N_k(t+\Delta t) - N_k(t) = \alpha \Delta t \frac{N_{k-1}(t) - N_k(t)}{N(0) + \alpha t}. \tag{5}$$

Note that the case of $k = 1$ is distinct. Each new node increments N_1 by one. So,

$$N_1(t+\Delta t) - N_1(t) = -\alpha \Delta t \frac{N_1(t)}{N(0) + \alpha t} + \alpha \Delta t. \tag{6}$$

These two equations can be condensed into one:

$$N_k(t+\Delta t) - N_k(t) = \frac{\alpha \Delta t}{N(0) + \alpha t}(N_{k-1} - N_k) + \alpha \Delta t \delta_{k,1}, \tag{7}$$

where $\delta_{k,1}$ is the Kronecker delta function (i.e., $\delta_{k,1} = 1$ if $k = 1$, and $\delta_{k,1} = 0$ otherwise). Dividing both sides by Δt and taking the limit $t \to 0$, we get the following differential equation for the dynamics of the expected degree distribution:

$$\dot{N_k} = \frac{\alpha}{N(0) + \alpha t}(N_{k-1} - N_k) + \alpha \delta_{k,1}, \tag{8}$$

where $\dot{N_k}$ is the first derivative of $N_k(t)$ with respect to time, and explicit dependence on time is omitted for expositional simplification.

To solve (8), we use the generating function $\psi(z,t) = \sum_k z^{-k} N_k(t)$, which is the conventional Z-transform in the k domain. Using (8) we get

$$\frac{\partial \psi(z,t)}{\partial t} = \frac{\alpha}{N(0) + \alpha t}(z^{-1} - 1)\psi(z,t) + \frac{\alpha}{z}. \tag{9}$$

So we intend to solve the following differential equation in the time domain:

$$\frac{\partial \psi(z,t)}{\partial t} - \frac{\alpha (z^{-1} - 1)}{N(0) + \alpha t}\psi(z,t) = \frac{\alpha}{z}. \tag{10}$$

This equation has the following general form:

$$\dot{\psi} + p(z,t)\psi = q(z,t), \tag{11}$$

whose solution is straightforward. Multiply both sides by an unknown integrating factor $\mu(z,t)$ to make both sides equal to $\frac{\partial}{\partial t}\big[\mu(z,t)\psi(z,t)\big]$. Then $\mu(z,t)$ is found to be $\exp[\int p(z,t)dt]$. Thus the final solution becomes:

$$\psi(z,t) = \frac{1}{\mu(z,t)}\bigg[\int \mu(z,t) q(z,t) dt + C(z)\bigg], \tag{12}$$

where C depends on the initial conditions. In our problem, the initial degree distribution is given and it will be used to determine C_z. For $\mu(z,t)$ we have:

$$\begin{aligned} \mu(t,z) &= \exp\bigg[-\int \frac{\alpha(z^{-1} - 1)}{N(0) + \alpha t} dt\bigg] \\ &= \exp\bigg[(1 - z^{-1}) \ln[N(0) + \alpha t]\bigg] \\ &= [N(0) + \alpha t]^{(1 - z^{-1})}. \end{aligned} \tag{13}$$

Using this and (12) , we find $\psi(z,t)$ as follows:

$$\psi(z,t) = \frac{\bigg[\int \frac{\alpha}{z}[N(0) + \alpha t]^{(1 - z^{-1})} dt + C_z\bigg]}{[N(0) + \alpha t]^{(1 - z^{-1})}}. \tag{14}$$

Note that the constant depends on z. Carrying out the integral, we arrive at:

$$\psi(z,t) = \frac{N(0) + \alpha t}{z(2 - \frac{1}{z})} + C_z (N(0) + \alpha t)^{\frac{1}{z} - 1}. \tag{15}$$

The constant C_z is obtained by looking at $t = 0$, when the following holds:

$$\psi(z,0) = \frac{N(0)}{z(2 - \frac{1}{z})} + C_z N(0)^{\frac{1}{z} - 1}. \tag{16}$$

Solving this for C_z and plugging the result in (15) yields

$$\begin{aligned} \psi(z,t) = \frac{N(0) + \alpha t}{z(2 - \frac{1}{z})} + \frac{N(0)}{N(0) + \alpha t}\bigg[\psi(z,0)(1 + \frac{\alpha t}{N(0)})^{\frac{1}{z}}\bigg] \\ - \frac{N(0)^2}{N(0) + \alpha t} \frac{1}{z(2 - \frac{1}{z})}\left(1 + \frac{\alpha t}{N(0)}\right)^{\frac{1}{z}}. \end{aligned} \tag{17}$$

To take the inverse transform, first note that:

$$\sum_{k=1}^{\infty} \frac{1}{2^k} z^{-k} = \frac{1}{z(2 - \frac{1}{z})}. \tag{18}$$

Also, denoting $\frac{\alpha t}{N(0)}$ by λ, note that we have:

$$(1 + \lambda)^{\frac{1}{z}} = e^{\frac{\ln(1+\lambda)}{z}}. \tag{19}$$

Using the Taylor expansion of the exponential, we get:

$$(1+\lambda)^{\frac{1}{z}} = \sum_{k=0}^{\infty} \frac{\left[\ln(1+\lambda)\right]^k}{k!} z^{-k}. \tag{20}$$

So the inverse transforms are:

$$\begin{cases} \dfrac{1}{z(2-\frac{1}{z})} \xrightarrow{\mathcal{Z}^{-1}} \dfrac{1}{2^k} u(k-1) \\ \\ \left(1+\dfrac{\alpha t}{N(0)}\right)^{\frac{1}{z}} \xrightarrow{\mathcal{Z}^{-1}} \dfrac{\left[\ln[1+\frac{\alpha t}{N(0)}]\right]^k}{k!}, \end{cases} \tag{21}$$

Where $u(x)$ is the Heaviside step function (i.e., $u(x) = 0$ for $x < 0$, and $u(x) = 1$ for $x \geq 0$). Finally, note that the multiplication of Z-transforms yields convolution after inversion. Let us denote the degree distribution of the initial graph by n_k, that is, n_k is the fraction of nodes at the outset with degree k. Thus inverting (17) gives:

$$\begin{aligned} N_k(t) &= \frac{N(0)+\alpha t}{2^k} u(k-1) \\ &+ \frac{N(0)^2}{N(0)+\alpha t} \left\{ n_k * \frac{\left[\ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!} \right\} \\ &- \frac{N(0)^2}{N(0)+\alpha t} \left\{ \left(\frac{u(k-1)}{2^k}\right) * \frac{\left[\ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!} \right\}, \end{aligned} \tag{22}$$

where $*$ denotes the convolution operator. To get the degree distribution, we divide this result by the total number of nodes at time t, which is equal to $N(0) + \alpha t$. So we obtain:

$$\begin{aligned} p_k(t) &= \frac{u(k-1)}{2^k} \\ &+ \left(\frac{N(0)}{N(0)+\alpha t}\right)^2 \left\{ n_k * \frac{\left[\ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!} \right\} \\ &- \left(\frac{N(0)}{N(0)+\alpha t}\right)^2 \left\{ \left(\frac{u(k-1)}{2^k}\right) * \frac{\left[\ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!} \right\}. \end{aligned} \tag{23}$$

As $t \to \infty$, the effect of initial conditions vanish. So the first term dominates. In this limit, the asymptotic degree distribution is:

$$\lim_{t\to\infty} p_k(t) = \frac{1}{2^k}. \tag{24}$$



FIG. 1: Degree distribution at time $t = 60$. The initial graph is 4−regular with $N(0) = 30$ nodes. Growth is under uniform probabilities and single attachments. At each timestep, one node is added, so $\alpha = 1$. The results are averaged over 20 Monte Carlo trials



FIG. 2: Growth under uniform probabilities and single attachments, on a ring of 50 nodes. The fractions of nodes having degree 3,4,5,6 are depicted in time. $\alpha = 1$ is used. The results are averaged over 20 Monte Carlo trials

Note that this matches the asymptotic behavior previously found for RRTs as presented in [38, 44, 45].

To compare theoretical prediction with simulation results, first we start off with a $4-$regular graph. This means that all nodes have 4 neighbors. We build this graph by first making a ring of 30 nodes, and then connect each node to two second closest neighbors. Fig. (1) shows the degree distribution at time $t = 60$, that is, $p_k(t = 60)$. The results are averaged over 20 Monte Carlo trials. Also, $\alpha = 1$, so nodes are introduced one at a time.

For the next simulation, we take a ring of 50 nodes and we plot $p_k(t)$ as a function of time, for $k = 3, 4, 5, 6$. The simulations and theoretical results are shown in Fig. (2).

B. Multiple Connections

Now, let us consider multiple attachments. Each new node that is introduced, chooses β existing nodes (where $\beta \geq 1$ is an integer) uniformly at random and links to them. An essential difference of this scheme from the previous one is that, if one starts from a disconnected graph, then the probability of ending up with a connected graph is nonzero. This probability was zero in the previous case, because each newly-introduced node only linked to one existing node and could not make a connection between two disconnected components.

Taking the similar steps that led to (7), the change in $N_k(t)$ is given by:

$$N_k(t+\Delta t)-N_k(t) = \frac{\beta\alpha\Delta t}{N(0)+\alpha t}(N_{k-1}-N_k)+\alpha\Delta t\delta_{k,\beta}. \tag{25}$$

Note that the last term indicates that each new node adds one to N_β, because its degree is β. Dividing by Δt, the differential equation for $N_k(t)$ becomes

$$\dot{N_k} = \frac{\beta\alpha}{N(0)+\alpha t}(N_{k-1}-N_k)+\alpha\delta_{k,\beta}\ . \tag{26}$$

Taking the Z-transform, (9) now takes the form:

$$\frac{\partial\psi(z,t)}{\partial t} = \frac{\alpha}{N(0)+\alpha t}(z^{-1}-1)\psi(z,t)+\frac{\alpha}{z^\beta}. \tag{27}$$

The following differential equation must be solved:

$$\frac{\partial\psi(z,t)}{\partial t} - \frac{\beta\alpha(z^{-1}-1)}{N(0)+\alpha t}\psi(z,t) = \frac{\alpha}{z^\beta}. \tag{28}$$

The integration factor is:

$$\begin{aligned}\mu(t,z) &= \exp\left[-\int\frac{\beta\alpha(z^{-1}-1)}{N(0)+\alpha t}dt\right]\\ &= \exp\left[\beta(1-z^{-1})\ln[N(0)+\alpha t]\right]\\ &= [N(0)+\alpha t]^{\beta(1-z^{-1})}.\end{aligned} \tag{29}$$

And from (12) , $\psi(z,t)$ is obtained:

$$\psi(z,t) = \frac{\left[\int\frac{\alpha}{z^\beta}[N(0)+\alpha t]^{\beta(1-z^{-1})}dt + C_z\right]}{[N(0)+\alpha t]^{\beta(1-z^{-1})}}. \tag{30}$$

After integration, we get:

$$\psi(z,t) = \frac{1}{z^\beta}\frac{N(0)+\alpha t}{1+\beta(1-z^{-1})} + \frac{C_z}{\left[N(0)+\alpha t\right]^{\beta(1-\frac{1}{z})}}. \tag{31}$$

Setting $t=0$, we find C_z:

$$C_z = \left[\psi(z,0) - \frac{N(0)z^{-\beta}}{1+\beta(1-z^{-1})}\right]N(0)^{\beta(1-z^{-1})}. \tag{32}$$

Substituting this C_z in (31) yields:

$$\begin{aligned}\psi(z,t) = {} & \frac{1}{z^\beta}\frac{N(0)+\alpha t}{1+\beta(1-z^{-1})}\\ & + \psi(z,0)\left[\frac{N(0)}{N(0)+\alpha t}\right]^{\beta(1-z^{-1})}\\ & - \frac{N(0)}{z^\beta\left[1+\beta(1-z^{-1})\right]}\left[\frac{N(0)}{N(0)+\alpha t}\right]^{\beta(1-z^{-1})}.\end{aligned} \tag{33}$$

Now we must invert this, term by term. First, note that:

$$\frac{1}{z^\beta\left[1+\beta(1-z^{-1})\right]} = (1+\beta)^{-1}\frac{1}{z^{\beta-1}\left[z-\frac{\beta}{\beta+1}\right]}. \tag{34}$$

Now, by definition, the inverse Z-transform is given by:

$$\begin{aligned}N_k(t) &= \frac{1}{2\pi i}\oint\psi(z,t)z^{k-1}dz\\ &= (1+\beta)^{-1}\oint\frac{z^{k-\beta}}{z-\frac{\beta}{\beta+1}}dz.\end{aligned} \tag{35}$$

In order to perform the integration, one must find the residuals of $\frac{z^{k-\beta}}{z-\frac{\beta}{\beta+1}}$ [46, 47]. For this purpose, we have to assume three distinct cases:

- If $k=\beta$, then we are looking for the residuals of $\frac{1}{z-\frac{\beta}{\beta+1}}$ which is unity.
- If $k>\beta$, then the only pole is at $z=\frac{\beta}{\beta+1}$ and the residual becomes $\left[\frac{\beta}{\beta+1}\right]^{k-\beta}$.
- If $k<\beta$, then $z=0$ is also a pole, and is of order $\beta-k$. The residual at this pole is equal to the following:

$$\frac{1}{(\beta-k-1)!}\frac{d^{(\beta-k-1)}}{dt^{(\beta-k-1)}}\frac{1}{z-\frac{\beta}{\beta+1}}\bigg|_{z=0} = -\left[\frac{\beta+1}{\beta}\right]^{\beta-k}. \tag{36}$$

The residual for the pole at $z=\frac{\beta}{\beta+1}$ is positive the same value, so they cancel out.

Putting these three together, we find the inverse transform:

$$\frac{1}{z^{\beta-1}\left[z-\frac{\beta}{\beta+1}\right]} \xrightarrow{\mathcal{Z}^{-1}} \left[\frac{\beta}{\beta+1}\right]^{k-\beta}u(k-\beta). \tag{37}$$

We have inverted the first term of (33). Another inverse transform that we need is the following.

$$\left[\frac{N(0)}{N(0)+\alpha t}\right]^{-\beta z^{-1}} \xrightarrow{\mathcal{Z}^{-1}} \frac{\left[\beta \ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!}. \tag{38}$$

Using these two results, the inverse transform of (33) becomes:

$$\begin{aligned} N_k(t) &= \frac{N(0)+\alpha t}{\beta}\left(\frac{\beta}{\beta+1}\right)^{k-\beta+1} u(k-\beta) \\ &+ \left[\frac{N(0)^{\beta+1}}{(N(0)+\alpha t)^{\beta}}\right]\left\{n_k * \frac{\left[\beta \ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!}\right\} \\ &- \left[\frac{N(0)^{\beta+1}}{\beta(N(0)+\alpha t)^{\beta}}\right] \times \\ &\left\{\left(\frac{\beta}{\beta+1}\right)^{k-\beta+1} u(k-\beta) * \frac{\left[\beta \ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!}\right\}, \end{aligned} \tag{39}$$

Then we divide by the total number of nodes $N(t) = N(0) + \alpha t$ to get the degree distribution:

$$\begin{aligned} p_k(t) &= \frac{1}{\beta}\left(\frac{\beta}{\beta+1}\right)^{k-\beta+1} u(k-\beta) \\ &+ \left(\frac{N(0)}{N(0)+\alpha t}\right)^{\beta+1}\left\{n_k * \frac{\left[\beta \ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!}\right\} \\ &- \left(\frac{N(0)}{N(0)+\alpha t}\right)^{\beta+1} \frac{1}{\beta} \times \\ &\left\{\left(\frac{\beta}{\beta+1}\right)^{k-\beta+1} u(k-\beta) * \frac{\left[\beta \ln\left(1+\frac{\alpha t}{N(0)}\right)\right]^k}{k!}\right\}. \end{aligned} \tag{40}$$

Now let us look at the long-time behavior of the result. When $t \to \infty$, the second and the third terms vanish. The first term prevails and tends to the following:

$$p_k(t) \sim \frac{1}{\beta}\left(\frac{\beta}{\beta+1}\right)^{k-\beta+1} u(k-\beta). \tag{41}$$

Note that for the case of $\beta = 1$, the same asymptotic distribution is obtained as the previous section. Also note that in the asymptotic limit, all nodes have degree at least β and the fraction of nodes with degree less than β tends to zero.



FIG. 3: Growth under uniform probabilities and multiple attachments with $\beta = 3$ and $\alpha = 1$, on a ring of 30 nodes. The fractions of nodes having degree 4,5,6,7 are depicted in time. The results are averaged over 30 Monte Carlo trials.



FIG. 4: Degree distribution at time $t = 30$ for a 6−regular graph, subject to growth with uniform probabilities and multiple attachments. The value of β is 3 and $\alpha = 1$. The results are averaged over 30 Monte Carlo trials

Fig. (3) shows the simulation results for a ring of 30 nodes, and $p_k(t)$ is depicted versus time, for $k = 4, 5, 6, 7$. Fig. (4) shows $p_k(t)$ for $t = 30$, for a 6−regular graph of total 30 nodes. The number of Monte Carlo trials is 30 for both cases. The value of β is 3 for both cases.

III. PREFERENTIAL LINKING

In this section we are going to focus on preferential attachment. New nodes, instead of selecting from the existing nodes uniformly at random, assign to them probabilities of connection, proportional to their degrees. So, each existing node has the chance of receiving a link from the newly-introduced node equal to its degree, divided by the sum of the degrees of every existing node. First we will assume the case where a new node only attaches to a single existing node, and then the case of multiple connections is considered.

A. Single Connection

As mentioned above, in the preferential attachment scheme, an existing node with degree k receives a link with probability $k/\sum_\ell \ell N_\ell$, where the denominator is the sum of the degrees of all existing nodes. So the probability that the destination node selected by a newly-born node has degree k is equal to $kN_k/\sum_\ell \ell N_\ell$. Using the same approach that led to (8), we arrive at the following differential equation for the evolution of $N_k(t)$:

$$\dot{N}_k = \frac{\alpha}{\sum_\ell \ell N_\ell}((k-1)N_{k-1} - kN_k) + \alpha\delta_{k,1}. \tag{42}$$

Now to proceed as before, we take the Z-transform of this equation. First note that if $X(z)$ is the Z-transform of a discrete function x_k, then $-z\frac{dX(z)}{dz}$ is the Z-transform for the the function kx_k. This means that, if the Z-transform of N_k is $\psi(z)$, then the Z-transform of the first term on the right hand side is as follows:

$$(k-1)N_{k-1} - kN_k \xrightarrow{\mathcal{Z}} (z-1)\frac{d\psi(z)}{dz}. \tag{43}$$

Second, note that the denominator of the attachment probabilities, $\sum_\ell \ell N_\ell$, is twice the number of links in the graph. Let us denote the number of links in the graph by $L(t)$. Note that, since each new node adds one new link, we have:

$$L(t) = L(0) + \alpha t \Longrightarrow 2L(t) = 2L(0) + 2\alpha t. \tag{44}$$

Twice the number of links in the initial graph equals $N(0)\bar{k}_0$, where $\bar{k}_0$ denotes the average degree of the initial graph. So we get:

$$2L(t) = N(0)\bar{k}_0 + 2\alpha t \Longrightarrow \sum_\ell \ell N_\ell = N(0)\bar{k}_0 + 2\alpha t. \tag{45}$$

We will temporarily use

$$\lambda \overset{\text{def}}{=} N(0)\bar{k}_0 \tag{46}$$

for brevity. The Z-transform of (42) is:

$$\frac{\partial\psi}{\partial t} - \frac{\alpha(z-1)}{\lambda + 2\alpha t}\frac{\partial\psi}{\partial z} = \alpha z^{-1}. \tag{47}$$

This is a first-order partial differential equation. We will solve this equation using the method of characteristics. For the convenience of the reader, we briefly shed light on how this method works through a simple example in the appendix. We refer the reader to [48–50], or other elementary references on partial differential equation, for further details.

We have the following system of equations:

$$\frac{dt}{1} = \frac{dz}{\left[-\frac{\alpha(z-1)}{\lambda+2\alpha t}\right]} = \frac{d\psi}{\alpha z^{-1}}. \tag{48}$$

The simplified version of the first equation is the following:

$$\frac{dt}{\lambda + 2\alpha t} = \frac{-dz}{\alpha(z-1)}. \tag{49}$$

Whose solution is the following:

$$(z-1)^2(\lambda + 2\alpha t) = C, \tag{50}$$

where C is a constant. The second equation is:

$$\frac{d\psi}{\alpha z^{-1}} = \frac{-dz}{\alpha(z-1)}(\lambda + 2\alpha t). \tag{51}$$

Replacing $(\lambda + 2\alpha t)$ by $C/(z-1)^2$ yields the following:

$$d\psi = \frac{-Cdz}{z(z-1)^3}. \tag{52}$$

Integrating both sides, gives:

$$\psi - C\left[\frac{-1}{z-1} + \frac{1}{2(z-1)^2} + \ln\frac{z}{z-1}\right] = C', \tag{53}$$

where C' is another constant. Following the lines of the example, we know that the solution has the following form for some function $\Phi(\cdot)$ which must be determined from the initial conditions:

$$\psi - C\left[\frac{-1}{z-1} + \frac{1}{2(z-1)^2} + \ln\frac{z}{z-1}\right] = \Phi\left[(z-1)^2(\lambda+2\alpha t)\right]. \tag{54}$$

Replacing C from (50) we get:

$$\begin{aligned}\Phi\big[(z-1)^2(\lambda + 2\alpha t)\big] &= \psi(z,t) \\ &- (\lambda + 2\alpha t)\left[-(z-1) + \frac{1}{2} + (z-1)^2\ln\frac{z}{z-1}\right].\end{aligned} \tag{55}$$

We must determine the function $\Phi(\cdot)$. Suppose the N_ks at the outset are known, so that for the initial graph we know $\psi(z,0)$. Let us denote it by $\psi_0(z)$. Setting $t = 0$ in (55) we get:

$$\begin{aligned}\Phi\big[(z-1)^2\lambda\big] &= \psi_0(z) \\ &- \lambda\left[-(z-1) + \frac{1}{2} + (z-1)^2\ln\frac{z}{z-1}\right].\end{aligned} \tag{56}$$

This helps us determine the function $\Phi(X)$. Denoting

$(z-1)^2\lambda$ by X, we get:

$$\Phi(X) = \psi_0(\sqrt{\frac{X}{\lambda}}+1)-\lambda\Big[-\sqrt{\frac{X}{\lambda}}+\frac{1}{2}+\frac{X}{\lambda}\ln\frac{\sqrt{X}+\sqrt{\lambda}}{\sqrt{\lambda}}\Big].$$

Since in (55) we have $[(z-1)^2\lambda]$ as X, let us explicitly find $\Phi((z-1)^2\lambda)$. We have:

$$\begin{aligned}&\Phi\big[(z-1)^2\lambda\big] = \psi_0\left[(z-1)\sqrt{\frac{\lambda+2\alpha t}{\lambda}}+1\right]\\&-\lambda\bigg[-(z-1)\sqrt{\frac{\lambda+2\alpha t}{\lambda}}+\frac{1}{2}\\&+(z-1)^2\frac{\lambda+2\alpha t}{\lambda}\ln\frac{(z-1)\sqrt{\lambda+2\alpha t}+\sqrt{\lambda}}{(z-1)\sqrt{\lambda+2\alpha t}}\bigg].\end{aligned} \tag{57}$$

We plug this expression in (55) and arrive at:

$$\begin{aligned}&\psi_0\left[(z-1)\sqrt{\frac{\lambda+2\alpha t}{\lambda}}+1\right]-\lambda\bigg[-(z-1)\sqrt{\frac{\lambda+2\alpha t}{\lambda}}\\&+\frac{1}{2}+(z-1)^2\frac{\lambda+2\alpha t}{\lambda}\ln\frac{(z-1)\sqrt{\lambda+2\alpha t}+\sqrt{\lambda}}{(z-1)\sqrt{\lambda+2\alpha t}}\bigg]\\&=\psi(z,t)-(\lambda+2\alpha t)\left[1-z+\frac{1}{2}+(z-1)^2\ln\frac{z}{z-1}\right].\end{aligned} \tag{58}$$

Now, let us make the following simplifications:

$$\begin{cases}\ln\dfrac{(z-1)\sqrt{\lambda+2\alpha t}+\sqrt{\lambda}}{(z-1)\sqrt{\lambda+2\alpha t}}=\ln\left[1+\dfrac{1}{z-1}\sqrt{\dfrac{\lambda}{\lambda+2\alpha t}}\right]\\ \ln\dfrac{z}{z-1}=-\ln(1-z^{-1}).\end{cases} \tag{59}$$

Using these simplifications and rearranging the terms in (58), we get:

$$\begin{aligned}&\psi(z,t)=\psi_0\left[(z-1)\sqrt{\frac{\lambda+2\alpha t}{\lambda}}+1\right]-2\alpha t(z-1)+\alpha t\\&-2\alpha 2(z-1)^2\ln(1-z^{-1})+(z-1)\lambda\sqrt{\frac{\lambda+2\alpha t}{\lambda}}\\&-\lambda(z-1)-\lambda(z-1)^2\ln(1-z^{-1})\\&-(z-1)^2(\lambda+2\alpha t)\ln\left[1+\frac{1}{z-1}\sqrt{\frac{\lambda}{\lambda+2\alpha t}}\right].\end{aligned} \tag{60}$$

Note that from the argument of the logarithm which was a result of integration, we know the region of convergence of the Z-transform is $z>1$, since the logarithm is not defined otherwise. Now let us define the new variable

$$c(t)\overset{\text{def}}{=}1-\sqrt{\frac{\lambda}{\lambda+2\alpha t}}. \tag{61}$$

This quantity is positive and less than unity at all times. Now, note that we have:

$$\begin{cases}1+\dfrac{1}{z-1}\sqrt{\dfrac{\lambda}{\lambda+2\alpha t}}=\dfrac{1-cz^{-1}}{1-z^{-1}}\\ (z-1)\sqrt{\dfrac{\lambda+2\alpha t}{\lambda}}+1=\dfrac{z-c}{1-c}\end{cases}. \tag{62}$$

So we simplify (60) further and arrive at:

$$\begin{aligned}&\psi(z,t)=\psi_0\left(\frac{z-c}{1-c}\right)-2\alpha t(z-1)+\alpha t\\&-(\lambda+2\alpha t)(z-1)^2\ln(1-z^{-1})+\lambda(z-1)\sqrt{\frac{\lambda+2\alpha t}{\lambda}}\\&-\lambda(z-1)-(\lambda+2\alpha t)(z-1)^2\ln\left(\frac{1-cz^{-1}}{1-z^{-1}}\right).\end{aligned} \tag{63}$$

Note that the two $\ln(1-z^{-1})$ terms cancel out. Also, note that there are three terms having the factor $(z-1)$. These three terms add up to:

$$\begin{aligned}&-2\alpha t(z-1)+\lambda(z-1)\sqrt{\frac{\lambda+2\alpha t}{\lambda}}-\lambda(z-1)\\&=(z-1)\sqrt{\lambda+2\alpha t}\Big(\sqrt{\lambda}-\sqrt{\lambda+2\alpha t}\Big)\\&=-(z-1)(\lambda+2\alpha t)c\end{aligned} \tag{64}$$

These simplifications transform (63) into the following:

$$\begin{aligned}\psi(z,t)=\psi_0\left(\frac{z-c}{1-c}\right)&+\alpha t-(z-1)(\lambda+2\alpha t)c\\&-(\lambda+2\alpha t)(z-1)^2\ln\left(1-cz^{-1}\right).\end{aligned} \tag{65}$$

Since $c\le 1$, $\forall t$ and $z>1$, we have $cz^{-1}<1$. So the logarithm can be Taylor-expanded. For $\ln(1-x)$ with $|x|<1$ we have:

$$\ln(1-x)\sim -x-\frac{x^2}{2}-\frac{x^3}{3}-\dots. \tag{66}$$

Using this, we have:

$$\begin{aligned}&(z-1)^2\ln(1-cz^{-1})\\&=\Big(-cz^{-1}-\frac{c^2z^{-2}}{2}-\frac{c^3z^{-3}}{3}-\dots\Big)\\&+\Big(-cz-\frac{c^2}{2}-\frac{c^3z^{-1}}{3}-\frac{c^4z^{-2}}{4}-\dots\Big)\\&+2\Big(c+\frac{c^2z^{-1}}{2}+\frac{c^3z^{-2}}{3}+\frac{c^4z^{-3}}{4}\dots\Big).\end{aligned} \tag{67}$$

Using this expression for the term with the logarithm in (65), we get:

$$\begin{aligned}\psi(z,t) &= \psi_0\left(\frac{z-c}{1-c}\right) + \alpha t - (z-1)(\lambda+2\alpha t)c \\ &+ (\lambda+2\alpha t)\bigg(cz^{-1} + \frac{c^2z^{-2}}{2} + \frac{c^3z^{-3}}{3} + \dots\bigg) \\ &+ (\lambda+2\alpha t)\bigg(cz + \frac{c^2}{2} + \frac{c^3z^{-1}}{3} + \frac{c^4z^{-2}}{4} + \dots\bigg) \\ &- 2(\lambda+2\alpha t)\bigg(c + \frac{c^2z^{-1}}{2} + \frac{c^3z^{-2}}{3} + \frac{c^4z^{-3}}{4}\dots\bigg).\end{aligned} \tag{68}$$

Note that this expression seemingly embodies terms with nonegative powers of z. Since the sequence of degree population N_k is zero for nonpositive values of k, the Z-transform is expected to only exhibit negative powers of z. So let us explicitly examine these terms and check that they do add up to zero. So let us rewrite the terms which embody nonnegative powers of z, which are:

$$\begin{aligned}&\alpha t - (z-1)(\lambda+2\alpha t)c + (\lambda+2\alpha t)cz \\ &+ (\lambda+2\alpha t)\frac{c^2}{2} - 2(\lambda+2\alpha t)c \\ &= \alpha t - (\lambda+2\alpha t)c + (\lambda+2\alpha t)\frac{c^2}{2}.\end{aligned} \tag{69}$$

The z^1 term readily vanishes. Now we focus on the constant terms. Using the fact that

$$(\lambda+2\alpha t)\frac{c^2}{2} = \lambda + \alpha t - \sqrt{\lambda}\sqrt{\lambda+2\alpha t}, \tag{70}$$

we have:

$$\begin{aligned}&\alpha t - (\lambda+2\alpha t)c + (\lambda+2\alpha t)\frac{c^2}{2} \\ &= -(\lambda+2\alpha t)c + \lambda + 2\alpha t - \sqrt{\lambda}\sqrt{\lambda+2\alpha t} \\ &= (\lambda+2\alpha t)\bigg(-c + 1 - \sqrt{\frac{\lambda}{\lambda+2\alpha t}}\bigg) \\ &= 0.\end{aligned} \tag{71}$$

So these terms do cancel out. Using this simplification, (68) becomes:

$$\begin{aligned}\psi(z,t) &= \psi_0\left(\frac{z-c}{1-c}\right) \\ &+ (\lambda+2\alpha t)\bigg(cz^{-1} + \frac{c^2z^{-2}}{2} + \frac{c^3z^{-3}}{3} + \dots\bigg) \\ &+ (\lambda+2\alpha t)\bigg(\frac{c^3z^{-1}}{3} + \frac{c^4z^{-2}}{4} + \frac{c^5z^{-3}}{5}\dots\bigg) \\ &- 2(\lambda+2\alpha t)\bigg(\frac{c^2z^{-1}}{2} + \frac{c^3z^{-2}}{3} + \frac{c^4z^{-3}}{4}\dots\bigg).\end{aligned} \tag{72}$$

which can be expressed in the following compact form:

$$\begin{aligned}\psi(z,t) &= \psi_0\left(\frac{z-c}{1-c}\right) \\ &+ (\lambda+2\alpha t)\sum_{k=1}^{\infty} z^{-k}\bigg(\frac{c^k}{k} - 2\frac{c^{k+1}}{k+1} + \frac{c^{k+2}}{k+2}\bigg)\end{aligned} \tag{73}$$

Now we have to take the inverse Z-transform. The sum on the right hand side is readily in the form of an expansion on z^{-1}. We focus on the first term on the right hand side. Let us denote the number of nodes in the initial graph who have degree k by $N_k(0)$. Then by definition, we have:

$$\psi_0(z) = \sum_k N_k(0)z^{-k}. \tag{74}$$

If we change the argument of ψ from z to $\frac{z-c}{1-c}$, we have:

$$\psi_0\left(\frac{z-c}{1-c}\right) = \sum_\ell N_\ell(0)\bigg(\frac{1-c}{z-c}\bigg)^\ell. \tag{75}$$

The inverse transform is given by:

$$\begin{aligned}&\frac{1}{2\pi i}\oint \sum_\ell N_\ell(0)\bigg(\frac{1-c}{z-c}\bigg)^\ell z^{k-1}dz \\ &= \sum_\ell N_\ell(0)\frac{(1-c)^\ell}{2\pi i}\oint \frac{z^{k-1}}{(z-c)^\ell}dz.\end{aligned} \tag{76}$$

Note that the residue of the function $\frac{f(z)}{(z-c)^\ell}$ for a differentiable function $f(\cdot)$, is given by $\frac{f^{(\ell-1)}}{(\ell-1)!}$, evaluated at $z=c$. For our problem, $f(z) = z^{k-1}$. So we have to evaluate the $(\ell-1)$-th derivative of the function z^{k-1}.

We have:

$$\frac{1}{(\ell-1)!}\frac{d^{(\ell-1)}}{dz^{(\ell-1)}}z^{k-1}\bigg|_{z=c}=\begin{cases}0 & \ell>k\\ \\ \dfrac{(k-1)!}{(\ell-1)!(k-\ell)!}c^{k-\ell} & \ell\leq k.\end{cases}\tag{77}$$

Using this, we find the integrals in (76) and arrive at:

$$\begin{aligned}&\frac{1}{2\pi i}\oint\sum_{\ell}N_{\ell}(0)\left(\frac{1-c}{z-c}\right)^{\ell}z^{k-1}dz\\&=\sum_{\ell}N_{\ell}(0)(1-c)^{\ell}c^{k-\ell}\frac{(k-1)!}{(k-\ell)!}.\end{aligned}\tag{78}$$

So the inverse Z-transform of (73) becomes:

$$\begin{aligned}N_k(t)=&\sum_{\ell}N_{\ell}(0)(1-c)^{\ell}c^{k-\ell}\binom{k-1}{\ell-1}\\&+(\lambda+2\alpha t)\left(\frac{c^k}{k}-2\frac{c^{k+1}}{k+1}+\frac{c^{k+2}}{k+2}\right).\end{aligned}\tag{79}$$

Replacing λ by $N(0)\bar{k}_0$, we get:

$$\begin{aligned}N_k(t)=&\sum_{\ell}N_{\ell}(0)\binom{k-1}{\ell-1}(1-c)^{\ell}c^{k-\ell}\\&+(N(0)\bar{k}_0+2\alpha t)\left(\frac{c^k}{k}-2\frac{c^{k+1}}{k+1}+\frac{c^{k+2}}{k+2}\right).\end{aligned}\tag{80}$$

Now to get the degree distribution, we divide this expression by the number of nodes at time t, which is equal to $N(0)+\alpha t$. As above, we denote the degree distribution of the initial graph by n_k, that is, n_k is the fraction of nodes at the outset with degree k. Thus the final result for the degree distribution is:

$$\begin{aligned}p_k(t)=&\frac{N(0)}{N(0)+\alpha t}\sum_{\ell}n_{\ell}\binom{k-1}{\ell-1}(1-c)^{\ell}c^{k-\ell}\\&+\frac{N(0)\bar{k}_0+2\alpha t}{N(0)+\alpha t}\left(\frac{c^k}{k}-2\frac{c^{k+1}}{k+1}+\frac{c^{k+2}}{k+2}\right),\end{aligned}\tag{81}$$

Combining (61) and (46), we get

$$c=1-\sqrt{\frac{N(0)\bar{k}_0}{N(0)\bar{k}_0+2\alpha t}}.\tag{82}$$



FIG. 5: Degree distribution at time $t=30$ for a 6−regular graph, subject to growth with preferential probabilities and single attachments. $\alpha=1$ and results are averaged over 30 Monte Carlo trials.

Note that as $t\to\infty$ we have:

$$\begin{cases}\lim\limits_{t\to\infty}\dfrac{N(0)}{N(0)+\alpha t}=0\\ \\ \lim\limits_{t\to\infty}\dfrac{N(0)\bar{k}_0+2\alpha t}{N(0)+\alpha t}=2\\ \\ \lim\limits_{t\to\infty}c=1.\end{cases}\tag{83}$$

Thus, the asymptotic behavior of the degree distribution is given by:

$$\lim_{t\to\infty}p_k(t)=2\left(\frac{1}{k+2}-\frac{2}{k+1}+\frac{1}{k}\right),\tag{84}$$

which simplifies to

$$\lim_{t\to\infty}p_k(t)=\frac{4}{k(k+1)(k+2)}.\tag{85}$$

As we mentioned previously, this asymptotic result was derived in [38, 39, 41]. Also, for large values of k, this pertains to the k^{-3} power law derived in [34].

Fig. (5) illustrates the simulation results and theoretical predictions for a ring of 30 nodes, and $p_k(t)$ is presented as a function of time, for $k=2,3,4$. Fig. (6) shows $p_k(t)$ for $t=30$, for a 6−regular graph of total 30 nodes.

B. Multiple Connection

Now let us consider the preferential attachment scheme again, but this time, each new node attaches to β existing nodes. This means that at time t, the number of nodes will be $N(0)+2\alpha t$. Also, at time t, the



FIG. 6: Growth under preferential probabilities and single attachments, on a ring of 30 nodes. The fractions of nodes having degree 2,3,4 are depicted in time. $\alpha = 1$ and results are averaged over 30 Monte Carlo trials.

sum of the degrees of all nodes (which equals twice the number of links) will be $N(0)\bar{k}_0 + 2\alpha\beta t$. Each newly born node adds one to N_β at that instant. Let us once again denote $N(0)\bar{k}_0$ by λ. Similar to (42), the evolution of $N_k(t)$ is

$$\dot{N}_k = \frac{\alpha\beta}{\sum_\ell \ell N_\ell}((k-1)N_{k-1} - kN_k) + \alpha\delta_{k,\beta}. \tag{86}$$

Taking the Z-transform leads us to:

$$\frac{\partial\psi}{\partial t} - \frac{\alpha\beta(z-1)}{\lambda + 2\alpha\beta t}\frac{\partial\psi}{\partial z} = \alpha z^{-\beta}. \tag{87}$$

Using the method of characteristics, we have :

$$\frac{dt}{1} = \frac{dz}{\left[-\frac{\alpha\beta(z-1)}{\lambda+2\alpha\beta t}\right]} = \frac{d\psi}{\alpha z^{-\beta}}. \tag{88}$$

The first equation yields a similar result as the previous section:

$$(z-1)^2(\lambda + 2\alpha\beta t) = C. \tag{89}$$

The second equation is the following:

$$\frac{d\psi}{\alpha z^{-\beta}} = \frac{-dz}{\alpha\beta(z-1)}\frac{C}{(z-1)^2}. \tag{90}$$

Let us define:

$$F(z) \stackrel{\text{def}}{=} \int^z \frac{x^{-\beta}}{(x-1)^3}dx. \tag{91}$$

Then we have:

$$\psi(z,t) = \frac{-C}{\beta}F(z) + \Phi(C). \tag{92}$$

This can be used to determine the unknown function $\Phi(\cdot)$. For $t = 0$ we have:

$$\psi_0(z) + \frac{(z-1)^2\lambda}{\beta}F(z) = \Phi\big[(z-1)^2\lambda\big]. \tag{93}$$

From this, we find that the function $\Phi(X)$ is

$$\Phi(X) = \psi_0\left(\sqrt{\frac{X}{\lambda}} + 1\right) + \frac{X}{\beta}F\left(\sqrt{\frac{X}{\lambda}} + 1\right). \tag{94}$$

Substituting C for X, (92) transforms to:

$$\psi(z,t) = \psi_0\left((z-1)\sqrt{\frac{\lambda + 2\alpha\beta t}{\lambda}} + 1\right) - \frac{(z-1)^2(\lambda + 2\alpha\beta t)}{\beta}\left[F(z) - F\left((z-1)\sqrt{\frac{\lambda + 2\alpha\beta t}{\lambda}} + 1\right)\right]. \tag{95}$$

Let us generalize (61) and define the following:

$$c \stackrel{\text{def}}{=} 1 - \sqrt{\frac{\lambda}{\lambda + 2\alpha\beta t}}. \tag{96}$$

As above, this quantity is less than one and tends to one as $t \to \infty$. Then (95) is simplified to:

$$\psi(z,t) = \psi_0\left(\frac{z-c}{1-c}\right) - \frac{(z-1)^2(\lambda + 2\alpha\beta t)}{\beta}\left[F(z) - F\left(\frac{z-c}{1-c}\right)\right]. \tag{97}$$

Now let us plug in the explicit form of $F(z)$. We found the result through speculation, by looking at the first few βs. The solution is:

$$F(z) = \int^z \frac{x^{-\beta}}{(x-1)^3}dx = \frac{\beta}{z-1} - \frac{1}{2(z-1)^2} + \frac{\beta(\beta+1)}{2}\ln(1 - z^{-1}) + \frac{1}{2}\sum_{k=1}^{\beta-1}\frac{(\beta-k)(\beta-k+1)}{k}z^{-k}. \tag{98}$$

Multiplying by $(z-1)^2$ as appeared in (97), we have:

$$G(z) \stackrel{\text{def}}{=} (z-1)^2F(z) = \beta(z-1) - \frac{1}{2} + \frac{\beta(\beta+1)}{2}(z-1)^2\ln(1 - z^{-1}) + \frac{1}{2}\sum_{k=1}^{\beta-1}\frac{(\beta-k)(\beta-k+1)}{k}(z^2 - 2z + 1)z^{-k}. \tag{99}$$

Now let us focus on taking the inverse Z-transform of

$G(z)$. Expanding the logarithm, we have:

$$\begin{aligned}G(z) &= \beta(z-1) - \frac{1}{2} \\ &+ \frac{\beta(\beta+1)}{2}(z-1)^2\left(-z^{-1} - \frac{z^{-2}}{2} - \frac{z^{-3}}{3} - \dots\right) \\ &+ \frac{1}{2}\sum_{k=1}^{\beta-1}\frac{(\beta-k)(\beta-k+1)}{k}(z^2-2z+1)z^{-k}. \quad (100)\end{aligned}$$

Note that the terms involving nonnegative powers f z must vanish, as above. First, let us look at the z^0 terms in $G(z)$. The second term gives:

$$\frac{\beta(\beta+1)}{2}\left(2 - \frac{1}{2}\right). \quad (101)$$

The third term in (100), gives two z^0 terms, one for $m = 1$ and one for $m = 2$. We have:

$$\frac{(\beta-2)(\beta-1)}{4} - \beta(\beta-1). \quad (102)$$

So the total z^0 term of $G(z)$ is:

$$-\beta - \frac{1}{2} + \frac{3\beta(\beta+1)}{4} + \frac{(\beta-2)(\beta-1)}{4} - \beta(\beta-1). \quad (103)$$

One can readily check that they do add up to zero.

Now let us examine the z^1 terms in $G(z)$. The second term in (100) gives:

$$\frac{\beta(\beta+1)}{2}(-1z). \quad (104)$$

The summation yields a z^1 term only for $m = 1$. This term is:

$$\frac{\beta(\beta-1)}{2}(z). \quad (105)$$

Plugging these in (100), we find that the total z^1 portion of $G(z)$ is:

$$\beta z + \frac{\beta(\beta+1)}{2}(-1z) + \frac{\beta(\beta-1)}{2}(z), \quad (106)$$

which adds up to zero.

To take the inverse Z-transform of (99), let us rewrite it as follows:

$$\begin{aligned}G(z) &= \frac{\beta(\beta+1)}{-2}\sum_{k=1}^{\infty}z^{-k}\left[\frac{1}{k} - \frac{2}{k+1} + \frac{1}{k+2}\right] \\ &+ \frac{1}{2}\sum_{k=1}^{\beta-1}\frac{(\beta-k)(\beta-k+1)}{k}(z^2-2z+1)z^{-k}. \quad (107)\end{aligned}$$

It can be further simplified as follows:

$$\begin{aligned}G(z) &= \frac{\beta(\beta+1)}{-2}\sum_{k=1}^{\infty}z^{-k}\frac{2}{k(k+1)(k+2)} \\ &+ \frac{1}{2}\sum_{k=1}^{\beta-1}z^{-k}\left\{\frac{(\beta-k)(\beta-k+1)}{k}\right. \\ &\left. - 2\frac{(\beta-k-1)(\beta-k)}{k+1} + \frac{(\beta-k-2)(\beta-k-1)}{(k+2)}\right\}. \\ &\qquad (108)\end{aligned}$$

Taking the common denominator of the terms inside the summation, this expression is simplified further and transforms to the following:

$$\begin{aligned}G(z) &= \frac{\beta(\beta+1)}{-2}\sum_{k=1}^{\infty}z^{-k}\frac{2}{k(k+1)(k+2)} \\ &+ \frac{1}{2}\sum_{k=1}^{\beta-1}z^{-k}\frac{2\beta(\beta+1)}{k(k+1)(k+2)} \quad (109)\end{aligned}$$

Since the terms are identical, $G(z)$ simplifies to the following compact form:

$$G(z) = -\sum_{k=1}^{\infty}\frac{\beta(\beta+1)}{k(k+1)(k+2)}z^{-k}. \quad (110)$$

Taking the inverse Z-transform is straightforward:

$$G(z) \xrightarrow{\mathcal{Z}^{-1}} \frac{-\beta(\beta+1)}{k(k+1)(k+2)}u(k-\beta). \quad (111)$$

Getting back to (97), we can rewrite it as follows:

$$\begin{aligned}\psi(z,t) &= \psi_0\left(\frac{z-c}{1-c}\right) \\ &- \frac{(\lambda+2\alpha\beta t)}{\beta}\left[G(z) - (1-c)^2 G\left(\frac{z-c}{1-c}\right)\right]. \quad (112)\end{aligned}$$

Note that we have:

$$(1-c)^2 = \frac{\lambda}{\lambda+2\alpha\beta t}, \quad (113)$$

so (112) is further simplified:

$$\begin{aligned}\psi(z,t) &= \psi_0\left(\frac{z-c}{1-c}\right) \\ &- \frac{(\lambda+2\alpha\beta t)}{\beta}\left[G(z) - \frac{\lambda}{\lambda+2\alpha\beta t}G\left(\frac{z-c}{1-c}\right)\right]. \quad (114)\end{aligned}$$

We have previously taken the inverse transform of the first term in (80), which was given by (78). Using the same procedure, we know how to extract the inverse transform of a function whose argument is $\left(\frac{z-c}{1-c}\right)$, once the inverse transform of the function is known. That is, knowing the inverse transform of $G(z)$, we can find the inverse transform of $G\left(\frac{z-c}{1-c}\right)$. Let us denote the inverse transform of $G(z)$ by g_k. Then we have:

$$G\left(\frac{z-c}{1-c}\right) \xrightarrow{\mathcal{Z}^{-1}} \sum_{\ell=1}^{k} g_\ell (1-c)^\ell c^{k-\ell} \binom{k-1}{\ell-1}. \tag{115}$$

Using this, along with (111), we take the inverse Z-transform of (114). Also, let us replace λ by $N(0)\bar{k}_0$. We wind up with:

$$\begin{aligned} N_k(t) &= \sum_{\ell=1}^{k} N_\ell(0)(1-c)^\ell c^{k-\ell}\binom{k-1}{\ell-1} \\ &+ \frac{(N(0)\bar{k}_0 + 2\alpha\beta t)}{\beta}\frac{\beta(\beta+1)}{k(k+1)(k+2)}u(k-\beta) \\ &- \frac{N(0)\bar{k}_0}{\beta}\sum_{\ell=\beta}^{k}\frac{\beta(\beta+1)}{\ell(\ell+1)(\ell+2)}(1-c)^\ell c^{k-\ell}\binom{k-1}{\ell-1} \end{aligned} \tag{116}$$

Finally, dividing (116) by the number of nodes at time t, which is equal to $N(0)+\alpha t$, yields the degree distribution at time t. As above, let us denote the degree distribution of the initial graph by n_k. The final result for the degree distribution is:

$$\begin{aligned} p_k(t) &= \frac{N(0)}{N(0)+\alpha t}\sum_{\ell} n_\ell(0)(1-c)^\ell c^{k-\ell}\binom{k-1}{\ell-1} \\ &+ \frac{(N(0)\bar{k}_0 + 2\alpha\beta t)}{N(0)+\alpha t}\frac{(\beta+1)}{k(k+1)(k+2)}u(k-\beta) \\ &- \frac{N(0)\bar{k}_0}{N(0)+\alpha t}\sum_{\ell=\beta}^{k}\frac{(\beta+1)}{\ell(\ell+1)(\ell+2)}(1-c)^\ell c^{k-\ell}\binom{k-1}{\ell-1} \end{aligned} \tag{117}$$

Now let us focus on the long time behavior, when $t \to \infty$, we have:

$$\begin{cases} \lim\limits_{t\to\infty} \dfrac{N(0)}{N(0)+\alpha t} = 0 \\ \lim\limits_{t\to\infty} \dfrac{N(0)\bar{k}_0 + 2\alpha\beta t}{N(0)+\alpha t} = 2\beta \\ \lim\limits_{t\to\infty} c = 1. \end{cases} \tag{118}$$



FIG. 7: Degree distribution at time $t = 30$ for a 6−regular graph, subject to growth with preferential probabilities and multiple attachments, with $\beta = 3$. The value of α is 1, and the results are averaged over 30 Monte Carlo trials.



FIG. 8: Growth under preferential probabilities and multiple attachments, on a ring of 30 nodes. The fractions of nodes having degree 2,3,4 are depicted in time. The value of β is 3 and the value of α is 1. The results are averaged over 30 Monte Carlo trials.

Using these values, the asymptotic degree distribution is obtained:

$$\lim_{t\to\infty} p_k(t) = \frac{2\beta(\beta+1)}{k(k+1)(k+2)}u(k-\beta), \tag{119}$$

which matches (2).

Fig. (7) is a depiction the simulation results and theoretical predictions for a ring of 30 nodes, and $p_k(t)$ is presented as a function of time, for $k = 2, 3, 4$. Fig. (8) shows $p_k(t)$ for $t = 50$, for a 6−regular graph of total 30 nodes. The value of β is 3 in both cases.

IV. SUMMARY AND FUTURE WORK

Previous work in the literature of network growth models mainly focus on the degree distribution of the graph in the asymptotic limit, that is, when the number of nodes tends to infinity and the effect of initial conditions can be neglected. In this contribution we found exact expressions for the expected degree distribution, which depend explicitly on the degree distribution of the initial graph. We considered two growth schemes. One in which new nodes choose from existing nodes uniformly at random, and then connect to them, and the other where these probabilities are proportional to degrees. Uniform and multiple attachments for the newly-born nodes are considered separately for both cases. Simulation results were accompanying theoretical predictions for each case.

One possible extension of the results presented in this work would be as follows. Suppose a given graph is subject to growth. The current state of the graph is known, and the growth mechanism can be approximated to be uniformly at random or be preferential attachment. Suppose quite on the contrary to the previous work in the literature, we are interested in the short-time behavior of the degree distribution. Then one could employ the results in this work, and expand the expressions in the vicinity of the initial condition up to arbitrary order of $(\alpha t)/N(0)$, and find the degree distribution perturbatively, to arbitrary precision.

Our analysis focuses on the expected degree distribution. $p_k(t)$ has a distribution of its own, whose mean value is presented in this work. One can also focus on the variance, or other statistical properties, of this distribution.

Appendix A: Method of Characteristics

Here we illustrate the basic procedure with an example. Consider the following partial differential equation for a function $\psi(x,y)$:

$$x^2\frac{\partial\psi(x,y)}{\partial x}+y^3\frac{\partial\psi(x,y)}{\partial y}=\psi^4(x,y)\quad ,x\geq 1, y\geq 0. \tag{A1}$$

First we solve the following system of equations:

$$\frac{dx}{x^2}=\frac{dy}{y^3}=\frac{d\psi}{\psi^4}. \tag{A2}$$

The first equation is solved as follows:

$$\frac{dx}{x^2}=\frac{dy}{y^3}\Longrightarrow\frac{1}{x}-\frac{1}{2y^2}=C_1, \tag{A3}$$

where C_1 is a constant. For the next equation we have (we arbitrarily pick one of the two remaining equations):

$$\frac{d\psi}{\psi^4}=\frac{dy}{y^3}\Longrightarrow\frac{1}{3\psi^3}-\frac{1}{2y^2}=C_2. \tag{A4}$$

Then the solution is of the form:

$$F(C_1)+G(C_2)=0, \tag{A5}$$

for any differentiable function $F(\cdot)$ and $G(\cdot)$. This is equivalent to:

$$C_2=H(C_1), \tag{A6}$$

for an arbitrary function $H(\cdot)$. This means that:

$$\frac{1}{3\psi^3}-\frac{1}{2y^2}=H\big(\frac{1}{x}-\frac{1}{2y^2}\big). \tag{A7}$$

This can be simplified to give:

$$\frac{1}{\psi^3}=\frac{3}{2y^2}+3H\big(\frac{1}{x}-\frac{1}{2y^2}\big), \tag{A8}$$

Denoting $3H(\cdot)$ by $\Phi(\cdot)$, the final solution can be expressed in the following general form for any differentiable function $\Phi(\cdot)$:

$$\psi(x,y)=\Big[\frac{3}{2y^2}+\Phi\big(\frac{1}{x}-\frac{1}{2y^2}\big)\Big]^{-\frac{1}{3}}. \tag{A9}$$

The function $\Phi(\cdot)$ is uniquely determined using the boundary conditions. Suppose we have the following information for the $x=1$ boundary:

$$\psi(1,y)=e^y. \tag{A10}$$

Plugging $x=1$ in (A9) yields:

$$\psi(1,y)=\Big[\frac{3}{2y^2}+\Phi\big(1-\frac{1}{2y^2}\big)\Big]^{-\frac{1}{3}}=e^y. \tag{A11}$$

Simplifying this, we get:

$$\Phi\big(1-\frac{1}{2y^2}\big)=e^{-3y}-\frac{3}{2y^2}. \tag{A12}$$

If we denote $1-\frac{1}{2y^2}$ by X, then $y=\sqrt{\frac{1}{2(1-X)}}$. So we arrive at:

$$\Phi(X)=\exp\Big[-3\sqrt{\frac{1}{2(1-X)}}\Big]-3(1-X). \tag{A13}$$

This $\Phi(\cdot)$ provides the unique solution of the form (A9), given the boundary condition $\psi(1,y)=e^y$. This concludes the example.

---

[1] M. Granovetter, “Strength of weak ties,” *American journal of Sociology*, vol. 79, pp. 1360–1380, 1973.

[2] S. Wasserman and K. Faust, *Social Network Analysis.* Cambridge: Cambridge University Press, 1994.

[3] A. Degenne and M. Forse, *Introducing Social Networks.* London: SAGE Publications Ltd., 1999.

[4] M. O. Jackson and B. Rogers, “The economics of small worlds,” *Journal of European Economic Association*, vol. 3, no. 617, 2005.

[5] V. Bala and S. Goyal, “A noncooperative model of network formation,” *Econometrica*, vol. 68, no. 5, p. 1181, 2000.

[6] L. A. Adamic, “The small world web,” *Research and Advanced Technology for Digital Libraries*, pp. 852–852, 1999.

[7] L. A. Adamic, R. M. Lukose, A. R. Puniyani, and B. A. Huberman, “Search in power-law networks,” *Physical review E*, vol. 64, no. 4, p. 046135, 2001.

[8] J. Leskovec, K. Lang, and M. Mahoney, “Empirical comparison of algorithms for network community detection,” in *Proceedings of the 19th international conference on World wide web.* New York, NY, USA: ACM, 2010, pp. 631–640.

[9] J. Leskovec, L. Adamic, and B. Huberman, “The dynamics of viral marketing,” *ACM Transactions on the Web (TWEB)*, vol. 1, no. 1, 2007.

[10] S. A. Delre, W. Jager, T. Bijmolt, and M. Janssen, “Will it spread or not? the effects of social influences and network topology on innovation diffusion,” *Journal of Product Innovation Management*, vol. 27, no. 2, pp. 267–282, 2010.

[11] B. J. Kim, T. Jun, J. Y. Kim, and M. Y. Choi, “Network marketing on a small-world network,” *Physica A*, vol. 360, 2006.

[12] A. C. R. Martins, C. de B. Pereira, and R. Vincente, “An opinion dynamics model for diffusion of innovations,” *Physica A*, vol. 388, pp. 3225–3232, 2009.

[13] M. E. J. Newman, “Spread of epidemic disease on networks,” *Physical Review E*, vol. 66, 2002.

[14] R. Pastor-Satorras and A. Vespignani, “Epidemic dynamics and endemic states in complex networks,” *Physical Review E*, vol. 63, 2001.

[15] M. Elowitz and S. Leibler, “A synthetic oscillatory network of transcriptional regulators,” *Nature*, vol. 4003, pp. 335—-338, 2000.

[16] S. Redner, “Citation statistics from 110 years of physical review,” *Physics Today*, vol. 58, no. 6, pp. 49–54, 2005.

[17] E. Ben-Naim, P. Krapivsky, and S. Redner, “Bifurcations and patterns in compromise processes,” *Physica D*, vol. 183, pp. 190–204, 2003.

[18] D. Acemoglu, A. Ozdaglar, and E. Yildiz, “Diffusion of innovations in social networks,” in *IEEE Conference on Decision and Control (CDC)*, 2011.

[19] P. Holme, C. Y. B.J. Kim, and S. K. Han, “Attack vulnerability of complex networks,” *Phys. Rev. E*, vol. 65, 2002.

[20] R. Cohen, K. Erez, D. ben Avraham, and S. Havlin, “Breakdown of the internet under intentional attack,” *Phys. Rev. Lett*, 2001.

[21] R. Pastor-Satorras and A. Vespignani, “Epidemic spreading in scale-free networks,” *Phys. Rev. Let.*, vol. 86, pp. 3200–3203, 2001.

[22] ——, “Immunization of complex networks,” *Physical Review E*, vol. 65, 2002.

[23] ——, “Epidemics and immunization in scale-free networks,” in *Handbook of Graphs and Networks: From the Genome to the Internet*, S. Bornholdt and H. Schuster, Eds., Wiley-VCH, Berlin, 2002, 2002.

[24] P. B. G. Rossmana, N. Esparzab, “I’d like to thank the academy, team spillovers, and network centrality,” *American Sociological Review*, vol. 75, no. 1, pp. 31–51, 2010.

[25] D.S.Price, “A general theory of bibliometric and other cumulative advantage processes,” *Journal of the American Society for Information Science*, vol. 27, no. 5, pp. 292–306, 2007.

[26] D. Price, “Statistical studies of networks of scientific papers,” in *Statistical Association Methods for Mechanized Documentation: Symposium Proceedings*, vol. 269. US Government Printing Office, 1965, p. 187.

[27] P. Erdős and A. Rényi, “On random graphs,” *Publicationes Mathematicae*, vol. 6, pp. 290–297, 1959.

[28] ——, “On the evolution of random graphs,” *Publications of the Mathematical Institute of the Hungarian Academy of Sciences*, vol. 5, pp. 17–61, 1960.

[29] B. Bollobás, *Random graphs.* Cambridge university press, 2001, vol. 73.

[30] ——, “The degree sequence of a scale-free random graph process,” *Random Structures and Algorithms*, vol. 18, no. 3, pp. 270–290, 2001.

[31] S. Watts and S. Strogatz, “The small world problem,” *Collective Dynamics of Small-World Networks*, vol. 393, pp. 440–442, 1998.

[32] M. Newman and D. Watts, “Renormalization group analysis of the small-world network model,” *Physics Letters A*, vol. 263, no. 4, pp. 341–346, 1999.

[33] A. Barabasi and R. Albert, “Emergence of scaling in random networks,” *science*, vol. 286, no. 5439, pp. 509–512, 1999.

[34] A. Barabasi, R. Albert, and H. Jeong, “Mean-field theory for scale-free random networks,” *Physica A: Statistical Mechanics and its Applications*, vol. 272, no. 1, pp. 173–187, 1999.

[35] H. Jeong, Z.Néda, and A. Barabasi, “Measuring preferential attachment in evolving networks,” *EPL (Europhysics Letters)*, vol. 61, no. 4, 2007.

[36] S. Zhou and R. Mondragón, “Accurately modeling the internet topology,” *Physical Review E*, vol. 70, no. 6, p. 066108, 2004.

[37] M. Newman, “Coauthorship networks and patterns of scientific collaboration,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, no. Suppl 1, pp. 5200–5205, 2004.

[38] P. Krapivsky, S. Redner, and F. Leyvraz, “Connectivity of growing random networks,” *Physical review letters*, vol. 85, no. 21, pp. 4629–4632, 2000.

[39] P. Krapivsky and S. Redner, "Organization of growing random networks," *Physical Review E*, vol. 63, no. 6, p. 066123, 2001.

[40] P. Krapivsky, G. Rodgers, and S. Redner, "Degree distributions of growing networks," *Physical Review Letters*, vol. 86, no. 23, pp. 5401–5404, 2001.

[41] P. Krapivsky and S. Redner, "Finiteness and fluctuations in growing networks," *Journal of Physics A: Mathematical and General*, vol. 35, no. 45, p. 9517, 2002.

[42] F.Chung, S. Handjani, and D. Jungreis, "Generalizations of polya's urn problem," *Annals of combinatorics*, vol. 7, no. 2, pp. 141–153, 2003.

[43] P. K. S. Redner, "Rate equation approach for growing networks," *Statistical mechanics of complex networks*, pp. 3–22, 2003.

[44] H. Na and A. Rapoport, "Distribution of nodes of a tree by degree," *Mathematical Biosciences*, vol. 6, pp. 313–329, 1970.

[45] S. Janson, "Asymptotic degree distribution in random recursive trees," *Random Structures & Algorithms*, vol. 26, no. 1-2, pp. 69–83, 2005.

[46] P. Morse and H. Feshbach, "Methods of theoretical physics, international series in pure and applied physics," *NY: McGraw Hill*, vol. 1, 1953.

[47] S. Hassani, *Mathematical Physics.* Springer, 1999.

[48] D. H. R. Courant, *Methods of Mathematical Physics.* John Wiley and Sons, 1989, vol. 2, ch. 2.

[49] E. Zauderer, *Partial differential equations of applied mathematics.* Wiley-Interscience, 2011, ch. 2.

[50] D. Zwillinger, *Handbook of differential equations.* Academic Press, 1998, ch. 2.